

## Expansion of the range of the Red-necked Wallaby in south-west Victoria

The Red-necked Wallaby *Macropus rufogriseus* is generally seen in woodlands where there is shrub cover, or on the fringes of forests. The species was thought to be substantially affected by past rabbit poisoning practices in woodlands and bushland reserves. It also appeared to be less common in its former haunts following the advent in the 1980s of the Black Wallaby *Wallabia bicolor*, which now occupies habitats from rocky ranges to wetland fringes or beach frontage. Visitors to Griffith Island at Port Fairy often see this species browsing on the introduced Shiny Leaf *Coprosma repens*, even wading in a pool to reach a favoured shrub.

The Black Wallaby was recognised for the first time in the Grampians in 1979 (Bird 1981). By 1987 it was seen in the Mount Napier State Park and at the Fulham Streamside Reserve on the Glenelg River near Balmoral. By 1992 it was a common sight in and around Hamilton (Bird 1992) and that situation continues, the wallabies exploiting any areas that provide dense cover (e.g. sightings in February 2011 at Lake Linlithgow and on the disused Hamilton-Coleraine Rail Reserve at Bochara). This species was previously unknown in the region, unlike the Red-necked Wallaby, which was often seen north of Hamilton (e.g. Grampians, Dundas Range, Fulham Streamside Reserve, Bear and Rocklands State Forests); west of Hamilton (e.g. Dergholm State Park, Hotspur State Forest, Weecurra State Forest); and, along with the Black Wallaby, south of Hamilton (e.g. Cobboboonee National Park, Lower Glenelg National Park, Pallisters Reserve near Orford, Homerton State Forest south of Mount Eccles National Park). Kay Aldridge (pers. comm.) from Heywood suggests that the Red-necked Wallaby is now common in the Cobboboonee forest, Homerton and Mount Clay, but tends to be encountered in small family groups rather than singly, as is usual for the Black Wallaby.

The first indication of a change in the range of the Red-necked Wallaby was the sighting of several animals on the summit of Mount Rouse,

the scoria cone at Peshurst, on 17 March 2007. This was most unexpected as the habitat seems to be atypical and the species was previously unknown there. The nearest bushland source is the Grampians, 25 km to the north. Unlike the Black Wallaby or Eastern Grey Kangaroo *Macropus giganteus*, both of which were also seen at Mount Rouse, the Red-necked Wallaby is not generally known to travel across open country far from shelter. How did they get to Mount Rouse?

The Red-necked Wallaby is recorded as a sub-fossil in bone deposits found in the Byaduk Caves (Wakefield 1964). The species was not recorded in surveys conducted from 1974 to 1995 in the contiguous volcanic landscape of Mount Napier State Park (Bird 1997), or on subsequent visits to the park up to 2010. It was a surprise therefore, to see three animals in the State Park on 16 January 2011, when driving along the Cole Track. One adult remained on the track for a minute or two, no more than 15 m distant from our vehicle but, at different sites, the others moved speedily into the dense Austral Bracken *Pteridium esculentum* in this open Manna Gum *Eucalyptus viminalis* forest. Where did these new arrivals come from?

One possibility is that the wallabies may have come from distant sites to Wildlife Shelters in the area. There are two shelters currently operating. Pam Turner (Wildwood Wildlife Shelter) on the Dunkeld-Moyston Road has had a marked increase in numbers of orphaned Red-necked Wallabies at her shelter (eight in the last twelve months), but fewer Black Wallabies. The wallabies are ultimately released in the Grampians by transporting them there or by 'soft release', where the animals eventually leave the premises when they choose to do so. Blue Gum plantations provide cover and presumably most animals re-enter the Grampians National Park or bushland adjacent to the park. However, it seems unlikely that the Mount Rouse animals were derived from that source, which is 35 km distant across open country.

Robyn Richardson runs a wildlife shelter at North Byaduk and uses a similar 'soft release' policy. Again, many more Red-necked Wallabies have been cared for in recent years, with a lesser number of Black Wallabies. The Red-necked Wallabies are mostly derived from the Heywood, Mount Eccles and Homerton area. In this case the released wallabies could easily find their way to the stoney rise areas of the Harmans Valley and Mount Napier State Park, only 3 km distant.

A second possibility is that the Red-necked Wallabies from forests 25 km west of Mount Napier or from Mount Napier to the south (I saw the species there on 21 May 2011) are using the cover of Blue Gum *Eucalyptus globulus* plantations, established in south-west Victoria from 1997 (Bird 2004), to migrate to new habitats, including Mount Napier. From 1997 to 2003, 100 000 ha of Blue Gums were planted on farms (Bird 2004), with a concentration in the higher rainfall areas such as Byaduk, Macarthur, Branxholme and Digby. By 2007 that total increased by about 30%, after which the major prospectus companies failed and the industry faltered. The planting provided corridors and 'stepping stones' that wildlife might use to migrate across a previously cleared landscape. Whether that could be the case for the Red-necked Wallaby population at Mount Rouse, 20 km east of Mount Napier, is doubtful,

since there are no plantations nearby and there is a much greater discontinuity of cover on the cleared farmland.

It seems probable that Red-necked Wallabies have been deliberately released at Mount Rouse, where there is sufficient cover from planted trees, shrubs, garden weeds and remnant Tussock Grass *Poa labillardierei* to sustain the small population. Whatever the cause, it will be interesting to see what areas they will colonise in future years, or if they can retain a presence in the 'newer volcanics' landscape of the Mount Napier State Park.

### References

- Bird PR (1981) A new macropod species of the Grampians. *The Victorian Naturalist* **98**, 67–68.  
 Bird PR (1992) Expansion of the range of the Black Wallaby in western Victoria. *The Victorian Naturalist* **109**, 89–91.  
 Bird PR (1997) Mammals and birds of the Mount Napier State Park. *The Victorian Naturalist* **114**, 52–66.  
 Bird, Rod (2004) Plantation and farm forestry developments in SW Victoria. *Hamilton Spectator*, 24 January, p. 23.  
 Wakefield NA (1964) Mammal sub-fossils from basalt caves in south western Victoria. *The Victorian Naturalist* **80**, 274–278.

### Rod Bird

21 Collins Street,  
 Hamilton, Victoria 3300  
 Email: rod.bird@bigpond.com

## Forty-three years Ago

### Friendly Rock Wallabies

These interesting notes are from Mr. A. G. Fellows of Alabama Hill, Charters Towers, North Queensland.

Over a period of twelve months we have won the trust of over thirty rock-wallabies that, instead of stampeding up our rocky hillside as once they did, now sit near our tool-shed door both in the early morning and late evening; or if not in sight, come bounding down just touching the rock tips when called by my wife or me as we break up stale bread loaves to throw to them. Soon the majority are sitting up straight, bread in paws, daintily though noisily munching away, heeding us very little, so that we often stand part surrounded by them, watching their antics. Occasional fights and chases occur between some individuals, whilst others make cat-like toilets after eating, and others drink daintily from the enamel water bowl which is often refilled for them.

From *The Victorian Naturalist* LXXXV, p. 90, April 4, 1968